

Long-term monitoring across elevational gradients (II): vascular plants on Pico Island (Azores) transect

COELHO, M.C.M., R.B. ELIAS, J. KLUGE, F. PEREIRA, D.S.G. HENRIQUES, S.C. ARANDA, P.A.V. BORGES, C. AH-PENG & R. GABRIEL



Coelho, M.C.M., R.B. Elias, J. Kluge, F. Pereira, D.S.G. Henriques, S.C. Aranda, P.A.V. Borges, C. Ah-Peng & R. Gabriel 2016. Long-term monitoring across elevational gradients (II): vascular plants on Pico Island (Azores) transect. *Arquipélago. Life and Marine Sciences* 33: 21 – 44.

Pico Island remains one of the last remnants of natural vegetation in the Azores, including the largest natural protected area; nevertheless, habitat change and the spread of exotic plants are visible, especially in those areas where human presence prevails. Currently, the lowlands are vastly occupied by pastures dominated by exotic herbs/grasses and most forests are dominated by *Pittosporum undulatum*. This paper aims to: i) review previous botanical studies related to elevational gradients; ii) investigate vascular plants composition and abundance in native vegetation, following an elevational transect (from 10 to 2200 m); and iii) investigate some patterns of the recorded diversity and distribution of vascular plants. Methodology follows a standardized protocol with observations in 100 m² plots. A total of 88 species were recorded, representing 35% of the indigenous but only 5% of the exotic species previously known from Pico. The richest areas were found between 600 and 1000 m and the areas with the lowest proportion of indigenous species occurred between 1800 and 2200 m. The recorded composition and richness values of endemic and native vascular plants support the high ecological and conservation value of the studied areas and constitute a good basis for long-term monitoring projects.

Key words: elevation, exotic species, indigenous vegetation, monitoring, vascular plants

Márcia C.M. Coelho (email: marcia.cm.coelho@uac.pt), Rui B. Elias, Fernando Pereira, Débora S.G. Henriques, Paulo A.V. Borges & Rosalina Gabriel, CE3C, Centre for Ecology, Evolution and Environmental Changes / Azorean Biodiversity Group and University of the Azores, Department of Agricultural Sciences, PT- 9700-042 Angra do Heroísmo, Portugal; and CITA-A and Portuguese Platform for Enhancing Ecological Research & Sustainability (PEERS). Jürgen Kluge, Faculty of Geography, Philipps University of Marburg, Deutschhausstrasse 10, DE- 35032 Marburg, Germany. Silvia C. Aranda, Museo Nacional de Ciencias Naturales, C/José Gutiérrez Abascal, 2, ES- 28006 Madrid, Spain. Claudine Ah-Peng, Université de La Réunion, UMR PVBMT, Pôle de Protection des Plantes, 7 Chemin de l'IRAT, 97410 Saint-Pierre, Ile de La Réunion, France.

INTRODUCTION

The Azores archipelago, in the middle North Atlantic, is formed by nine isolated islands, colonized since the 15th century and currently bearing a small fraction of their original primary forest habitats (estimated at less than 3%; Triantis et al. 2010; Gaspar et al. 2011). Even if about a

fifth (21%) of the Azores terrestrial territory is currently under some level of protection, native forest communities are restricted to much smaller fragments, more consentaneous with the areas categorized as natural reserves (8%) and the islands of Terceira, Pico and Flores are the largest contributors to that percentage (PNA [cited 2016]).

In the Azores, the known vascular flora comprises 1110 species and subspecies: 881 exotic (79%) (including 328 casuals); 73 endemic (7%) and 136 (12%) native non-endemic plants to the archipelago (Silva et al. 2010); the remaining 20 species have a doubtful/undetermined origin. The proportion of endemic vascular plants in the Azores is low, if compared with other archipelagos of Macaronesia, as Madeira (154/1204) (Borges et al. 2008) and Canary Islands (539/2091) (Arechavaleta et al. 2010), but relatively high if compared with most European regions, such as mainland Portugal (150/3314) (Menezes de Sequeira et al. 2012). If only indigenous species are considered, the endemism rate of Azorean flora, reaches about one third (35%), a value higher than in Madeira (20%). Moreover, there is a recent recognition that current number of endemic angiosperms in Azores is underestimated (Schäfer et al. 2011). Within this context, of high level of both exotic/invasive species, and endemic biodiversity, it is important to monitor the remaining fragments of native vegetation, to be able to evaluate the trends of habitat and climate change, invasion by exotics and the success of conservation programs (Gabriel et al. 2014).

CHARACTERIZATION OF PICO ISLAND

Pico Island is located between the coordinates 38°33'57" and 38°33'44" N of latitude and 28°01'39" and 28°32'33" W of longitude (França et al. 2003). With a surface of 445 km² and a coastal perimeter of 126 km (Forjaz 2004), it is the largest of the five islands of the Central Group. Pico is shaped roughly as an oval, elongated in the WNW–ESE axis (45 to 50 km from “Madalena” to “Ponta da Ilha”), with a maximum width in the NNE–SSW axis (16 km from “Arcos” to “Ponta de S. Mateus”) (Zbyszewski et al. 1963).

This is the youngest of the nine Azorean islands, holding the highest mountain of Portugal (2351 m a.s.l.), an active stratovolcano that created the island 250 000 years ago (França et al. 2003). Its cone rises abruptly from the sea, with a wide base that narrows from 1200 m elevation and terminates in a shallow caldera (c. 3 km wide), from the centre of which a secondary cone,

called “Piquinho” rises over 60 m (Forjaz 2004, França et al. 2014). Due to its volcanic origin, Pico Island is mostly formed by basaltic rocks (basalts, mugearite, hawaiiite) and pyroclastic materials of basaltic composition (Madeira et al. 2007). Unlike other Azorean islands, such as São Jorge, the coastal areas have almost no high cliffs, and are characterized by the low lava flows and a few sandy beaches of basaltic origin (Nunes 1999).

Pico Island is quite near its neighbouring islands, Faial and São Jorge, separated only by two narrow channels, the first with 6 km width and 95 m of depth, and the second with 18 km width and 1200 m of depth (Nunes 1999). During the last glacial maximum (20 000 years ago) Pico and Faial Islands were connected (Rijsdijk et al. 2014) which may have been of great biological value allowing the exchange of several species.

Pico Island has a very low urban density (32 inhabitants/km²) compared with the islands of Terceira (141 inhabitants/km²) and São Miguel (185 inhabitants/km²) (INE [cited 2015]).

Pastures for dairy cattle (40%) and exotic forests (33%) of *Cryptomeria japonica*, *Pinus pinaster* and *Acacia melanoxylon* dominate the land use of Pico Island (Cruz et al. 2007), while agriculture (8%; including viticulture and maize), is much less predominant. The economy of the island is mainly based on livestock (with an important production of cheese), fishery, agriculture (mainly winery), and tourism (Lima et al. 2014; Silva 2015).

The climate of Pico Island is considered humid mesothermal with oceanic characteristics. Annual precipitation varies between 974 mm at low elevations and 4834 mm at 1200 m, and decreases for 2025 mm at summit (Azevedo 1996; Azevedo et al. 1999a,b). The average air temperature of February (the coldest month) ranges from 15° C (in the coastal areas) to 6° C (in the top of the mountain) and in August (the warmest month) from 23° C to 13° C, respectively (Azevedo et al. 2004; Barceló & Nunes 2011). This island is the only that presents snowfall in its higher lands, persisting for more or less extended periods of time, during the winter months (Azevedo et al. 2004). The annual relative air humidity in Pico is below 80% along the coastal areas, increasing in

the remaining elevations (80–100%) (Azevedo et al. 2004). However, exceptional conditions of air dryness occur, with the relative humidity reaching values lower than 10% at the summit of Pico Mountain (França et al. 2014). The annual average wind speed in coastal areas, during winter months, reaches 20 km/h but gusts commonly attain speeds of 100 km/h; in summer months the wind velocity values decrease to under 10 km/h (França et al. 2014).

Pico's Natural Park is the largest of the archipelago, covering more than one third of the island's area, and aiming to shelter the endemic fauna and flora of the Azores, including their unique subalpine and alpine habitats (Leuschner 1996). However, land categorized as Strict Nature Reserve (IUCN category Ia) covers only 5.2% of the island, including the mountain of Pico, "Caveiro", "Mistério da Prainha" and "Furnas de Santo António" (DLR 20/2008/A).

The island hosts more than four fifths of all indigenous vascular plants found in the Azores (N=173; 83%) (Silva et al. 2010) but, among its 616 *taxa*, it also includes 430 exotic *taxa* (naturalized and casual), that are considered introduced and 13 *taxa* with doubtful colonization status (Borges et al. 2010; Silva et al. 2010). The invasive *Pittosporum undulatum* Vent. has been occupying large areas of Azorean forest, about a third of all forest surface of the nine islands (Silva et al. 2008c), but that increase is more prominent in Pico Island since that species occupies 61% of all forest surface (REAA 2014). However, the climatic optimum of this species is at mid-altitudes being unable to live in the colder and much more humid altitudes (Hortal et al. 2010).

VEGETATION ZONATION STUDIES IN PICO ISLAND

Since the settlement of Portuguese in the Azores, the vegetation suffered significant alterations, mainly due to land-use change and the introduction of exotic species. In the 16th century, Gaspar Frutuoso (1998), in the series "Saudades da Terra" described the Azorean vegetation as composed of "tall and dense forests", without mentioning any particular stratification according to elevation. Among the woody species assigned to Pico Island by that author, it is possible to recognize *Juniperus brevifolia* (Seub.) Antoine

("Zimbro, Cedro"/Azorean cedar), *Taxus baccata* L. ("Teixo"/Yew), *Erica azorica* Hochst. ex Seub. ("Urze"/Azorean Heather), *Frangula azorica* V. Grubov ("Sanguinho"/Buckthorn), *Laurus azorica* (Seub.) Franco ("Louro"/Azorean sweet bay), *Morella faya* (Aiton) Wilbur ("Faia"/Firetree), *Myrsine africana* L. ("Tamujo"/Cape myrtle), *Picconia azorica* (Tutin) Knobl. ("Pau-branco"), and *Viburnum treleasei* Gand. ("Folhado"/Azorean laurustinus) (Frutuoso 1998).

Much later, several botanists attempted to describe the Azorean vegetation. Many of these early endeavours used different elevation belts and focused on the highest island of the archipelago (*cf.* Table 1). In fact, its wide range of elevations (more than twice the elevation of the second highest point in the Azores — São Miguel, 1105 m elevation [Forjaz 2004]) allows the presence of more vegetation belts and comparisons with other mountain regions, either in Macaronesia or elsewhere.

Table 1, adapted and updated from Lüpnitz (1975), summarizes the works of previous authors that explicitly used elevation as an important descriptive variable in the delimitation of vegetation belts in the Azores, such as Seubert & Hochstetter (1843), Morelet (1860), Guppy (1917), Allorge & Allorge (1946), Marler & Boatman (1952), Tutin (1953), Dansereau (1966), Lüpnitz (1975), Purvis et al. (1994, 1995). However, other very important works, namely by Sjögren (1973, 1978), Dias (1996), Elias (2001, 2007), Dias et al. (2004) and Elias et al. (2011) followed a different approach to describe Azorean plant communities, the first by using a phytosociological perspective, the others a more quantitative multivariate analysis, focusing on the communities without restricting their analysis to the finding of elevation belts.

Since the pioneer work by Seubert & Hochstetter (1843), more than 150 years separate these visions of the Azorean vegetation, and this was a dynamic period regarding land use (*e.g.* Triantis et al. 2010; Silveira 2013), with massive loss of native habitats.

Considering Table 1 (in the following page), it is possible to recognize some consensus towards a vegetation change around 500–600 m and another around 1500 m, but the descriptions of the

Table 1. Summary of different systems attempting to describe Azorean vegetation, using different elevational belts (adapted and updated from Lüpnitz 1975; see details in text)

Elevational belt (m)	Seubert & Hochstetter (1843)	Morelet (1860)	Guppy (1917)	Allorge & Allorge (1946)	Marler & Boatman (1952)	Tutin (1953)	Dansereau (1970)	Purvis et al. (1994:upper elevations; 1995: lower elevations)		
1-100	Cultivated or Mediterranean zone (0 m - 500 m)	Zone of cultivation (0 m - 500 m)	Lower woods or Faya zone (0 m – 610 [800] m)	Littoral, Mediterranean or Riccia region (0 m - 200 [300] m)	Rocky slopes (0 m - 50 [100] m)		Coastal vegetation (0 m - 100 m)			
100-200					Cleared woodland with <i>Pinus pinaster</i> (50 m - 150 m)					
200-300					<i>Pittosporum undulatum</i> woodland (150 m - 500 m)				Laurel forest (broad-leaved evergreen trees) (100 m - 600 m)	<i>Morella faya</i> (0 m - 300 m)
300-400										
400-500										<i>Laurus azorica</i> (300 m – 650 m)
500-600	Canarian zone or belt of laurel wood (500 m - 850 m)	Zone of woods: laurels and other evergreen trees and shrubs (500 m - 1500 m)	Upper woods or <i>Juniper</i> zone (610 m - 1237 m)	<i>Sphagnum</i> region (<i>Juniperus brevifolia</i> and <i>Persea indica</i> forests) (600 m -700 [800] m)	<i>Myrica faya</i> and <i>Persea azorica</i> (500 m - 1000 m)		<i>Ericetum azoricae</i> (<i>Erica azorica</i> and <i>Juniperus brevifolia</i>) (600 m - 1500 m)	<i>Juniperus brevifolia</i> (450 m – 1100 m)		Mid-elevation lichen-rich forest (700 m - 900 m)
600-700										
700-800										
800-900										
900-1000	Azorean zone or region of shrubs (850 m - 1500 m)	Zone of woods: laurels and other evergreen trees and shrubs (500 m - 1500 m)	Shrubs or <i>Juniper</i> zone (1237 m - 1676 m)	<i>Juniperus brevifolia</i> region (<i>Adelanthus decipiens</i> and <i>Daltonia stenophylla</i>) (800 [900] m - 1400 [1500] m)	Calluna vulgaris heath grasslands and wet hollows (1000 m - 1500 m)	Mixed <i>Erica azorica</i> – <i>Juniperus brevifolia</i> scrub (1000 m - 1200 m)			<i>Erica azorica</i> (650 m-1600 m)	
1000-1100										
1100-1200										
1200-1300							<i>Juniperion brevifolii</i> (1200m-1300m)			
1300-1400								<i>Calluna vulgaris</i> - <i>Daboecia azorica</i> heath land; <i>Erica</i> - sparse (1340 m - 1500 m)		
1400-1500	Bush or scrub zone (1500 m - 1700 m)	Zone of woods: laurels and other evergreen trees and shrubs (500 m - 1500 m)	Shrubs or <i>Juniper</i> zone (1237 m - 1676 m)					Upper elevation lichen-rich forest (1050 m - 1150 m)		
1500-1600									Upper zone: more pastures and heaths (1500 m - 2351 m)	<i>Calluna, Menziesia</i> and <i>Thymus</i> zone (1676 m - 2351 m)
1600-1700	Summit zone (1700 m - 2351 m)						At least 50 % of bare rock (2300 m - 2351 m)			
1700-1800										
1800-1900										
1900-2000										
2000-2100										
2100-2200										
2200-2300										
2300-2351										

communities differ considerably among the authors. Nevertheless, Guppy (1917), Allorge & Allorge (1946), Tutin (1953), Dansereau (1966) and Lüpnitz (1975) separate the lower elevation vegetation belts, dominated by broadleaf species (generally below 600 m), from the higher elevation *Juniperus* zone.

From 600 to 1500 m, Tutin (1953) described *Ericetum azoricae*. However, it is the endemic conifer *Juniperus brevifolia* that is generally recognized as the main architect species in the Azores montane vegetation. Guppy (1917) described forests of *Juniperus brevifolia* and *Laurus azorica* (832–1485 m), while Allorge & Allorge (1946) mentioned pure stands of *Juniperus brevifolia* between 850 and 1500 m. Sjögren (1973), divided the plant communities of the Azores into four alliances, defining the broad alliance *Juniperion* around the presence of *Juniperus brevifolia*; the importance of this gymnosperm is lately emphasized by Elias (2007).

It seems to be consensual that *Calluna vulgaris* (L.) Hull and *Thymus caespititius* Brot. dominate the vegetation above 1500 m, but Purvis and colleagues (1994, 1995), working with lichens, were able to describe a zonation, even above that elevation. It is worth mentioning that the *Callunetum* put forward by Tutin (1953) was later questioned by Sjögren (1973), who proposed an impoverished *Juniperion* for all the mountainous area of Pico.

The strong influence of the human presence was already noted in the 19th century by Seubert & Hochstetter (1843), which referred an “area of cultures” to the elevations below 500 m. That same idea is reinforced by Marler & Boatman (1952), who introduced *Pittosporum undulatum* as a differential species of *Pittosporum*, a phytosociological association, claiming that “the vegetation of the northern slopes of Pico has been affected by the recently assumed dominance of an introduced tree *Pittosporum undulatum*”, with the consequent disappearance of natural vegetation (1952, p. 143–144).

This is the second paper on the series “Long-term monitoring across elevational gradients” that aims to identify and characterize the biodiversity of native flora in several Azorean islands along an elevational transect. This paper presents the results of the MOVECLIM field study concerning

the vascular flora of Pico Island (further details in Gabriel et al. 2014). The main goals of this work are to i) review previous botanical studies related to elevational gradients — focusing on Pico Island, since it has the widest elevational range; ii) investigate vascular plants composition and abundance in key areas of native vegetation, following an elevational transect (from 10 to 2200 m); and iii) investigate some patterns of the recorded diversity and distribution of vascular plants.

METHODS

STUDY SITE AND SAMPLING DATES

Up to 1000 m, the Pico Island’s transect was set up along the eastern side of the island (Fig. 1a) in order to encompass the best preserved native vegetation and avoiding as much as possible areas with exotic species. From 1200 to 2200 m the transect roughly followed the mountain trail on the south declivity of the main volcanic cone (Fig. 1b). All sites, except the one studied at 400 m, were placed inside the limits of the island’s Natural Park, namely in Nature Reserves areas as Pico mountain, “Caveiro” and “Mistério da Prainha”. There were three field study phases: 1st) 4–12 September 2012, with the purpose to setting the permanent plots in 12 sites, along an elevational gradient, from 10 to 2200 m, using a 200 m elevational step (Fig. 1) and conduct a preliminary survey of the vascular species present in each of the plots; 2nd) 5–11 May 2013, the vascular species cover was obtained on the same plots; and 3rd) 28 July to 4 August 2013, the sites were revisited to gather pteridophyte data.

SAMPLING METHOD

The sampling methodology mainly followed the standardized protocol BRYOLAT (Kessler et al. 2011; Ah-Peng et al. 2012), and was thoroughly described in the first paper of this series (Gabriel et al. 2014). This methodology has been designed for the study of bryophyte and fern diversity and distribution along environmental gradients in the tropics and subtropics, but it also allows the characterization of vascular flora by recording the presence and cover, as well as the heights and diameter at breast height (dbh) of the largest trees.



Fig. 1. Location of Pico Island (a) and of the transect, showing 12 sites, sampled with 200 m elevational steps (b, c). (b) Lower elevation sites, from 10 to 1000 m and (c) Mountainous sites, from 1200 to 2200 m (surveys in 2012 and 2013).

Vascular plants were identified and cover percentage visually estimated in two 100m² plots per each site (species identified to genera did not have reproductive structures available in the field to assist in further identification). The taxonomical classification of the vascular flora follows Silva et al. (2010).

CLIMATE DATA IN THE MOVECLIM TRANSECT

Climate data was obtained from the CIELO Model (Azevedo 1996, Azevedo et al. 1999a,b). CIELO is a physically based model that simulates the climatic variables on an island using data from the synoptic reference of a meteorological site. For this work, the used variables were: precipitation (total annual), relative humidity (maximum and minimum) and temperature (maximum and minimum), obtained from an estimate of the values from the previous 30 years (Table 2).

DATA ANALYSIS

Species richness and cover–abundance were analysed for each plot. The occurrence of a Mid Domain Effect (MDE) (Colwell & Hurtt 1994; Colwell & Lees 2000) was tested, using indigenous and total richness of vascular plants. When the MDE is confirmed, species distribution ranges overlap more towards the middle of the elevational gradient, due to topographical and/or geographical constraints.

The disturbance index (D) (Cardoso et al. 2013) was obtained to determinate the landscape disturbance accounting specifically for landscape composition and configuration. This index reflects an anthropogenic disturbance gradient, ranging from zero (no disturbance at all) to 100 (maximum possible disturbance, like an urban/industrial area) (Table 2).

Vascular plants on Pico Island transect

Table 2. Geographic data, climate data [Rain (total annual); Relative humidity (average annual) and Temperature (average annual) (Azevedo 1996; Azevedo et al. 1999a,b)], biological data and disturbance index [minimum 0; maximum 100 (Cardoso et al. 2013)], for the 12 sites (24 plots) in Pico Island. Site codes: standard elevation (a.s.l.) and plot number.

Site name	Site codes	Geographic data					Climate data			Biological data				
		Coordinates (N)	Coordinates (W)	Elevation (m)	Slope (°)	Exposure (°)	Rain (mm)	Relative humidity (%)	Temperature (° C)	Endemic AZ	Endemic MAC	Native	Naturalized	Doubtful
1."Manhenha"	0010P1	38°24'49.5"	28°01'47.3"	17	5	205	974	86	19	6	4	4	38	
	0010P2	38°24'49.5"	28°01'47.8"	14	10	300				5	3	6		
2."Cabeço da Hera"	0200P1	38°25'05.5"	28°03'14.2"	224	50	270	1014	93	17	3	5	6	47	
	0200P2	38°25'05.4"	28°03'13.1"	226	50	60				3	6	8		
3."Fetais, Piedade"	0400P1	38°25'32.8"	28°05'15.5"	365	5	140	1292	94	16	10	4	3	57	
	0400P2	38°25'33.3"	28°05'14.5"	364	5	25				8	5	4		
4."Caminho dos Burros, Chão Verde"	0600P1	38°28'7.85"	28°16'34.3"	621	10	180	3066	96	14	19	1	15	4	15
	0600P2	38°28'7.73"	28°16'33.4"	623	15	225				18	1	15	4	
5."Caiado"	0800P1	38°27'20.8"	28°15'26.2"	809	5	320	3513	98	13	12	1	9	1	27
	0800P2	38°27'20.4"	28°15'26.1"	813	5	110				13	1	8	2	
6."Caveiro"	1000P1	38°26'13.8"	28°12'46.9"	952	25	20	3812	99	12	14	1	7	2	23
	1000P2	38°26'13.8"	28°12'46.1"	947	25	350				17	2	9		
7.Pico Mountain	1200P1	38°28'13.3"	28°25'30.9"	1261	5	320	4834	98	11	11	9	6	2	25
	1200P2	38°28'14.3"	28°25'30.1"	1261	5	300				11	7		1	
8.Pico Mountain	1400P1	38°28'10.5"	28°25'16.9"	1418	20	70	4653	98	10	8	4		1	24
	1400P2	38°28'09.5"	28°25'16.5"	1407	15	95				8	4	2	1	
9.Pico Mountain	1600P1	38°27'57.5"	28°24'59.5"	1601	30	95	4047	98	9	9	8	2		24
	1600P2	38°27'55.7"	28°24'59.7"	1588	25	100				10	8	2		
10.Pico Mountain	1800P1	38°27'57.9"	28°24'45.2"	1805	35	70	3613	98	8	3	3			24
	1800P2	38°27'57.4"	28°24'45.0"	1802	35	70				3	3			
11.Pico Mountain	2000P1	38°27'56.6"	28°24'30.0"	2003	40	90	2723	99	7	1	2			23
	2000P2	38°27'56.1"	28°24'29.2"	2009	40	120				1	2			
12.Pico Mountain	2200P1	38°27'58.9"	28°23'57.2"	2239	10	320	2025	99	6	1	2			23
	2200P2	38°28'00.2"	28°23'57.9"	2242	15	320				1	2			

RESULTS

PLOT DESCRIPTION

Twelve elevation sites were sampled on Pico Island, comprising 24 plots (Fig. 1, 2 and 3; Table 2). Climate becomes harsher along the elevational gradient. Around 1600 m there is a clear change of conditions (inversion layer), where precipitation values decrease (Table 2).

The list of vascular plant species registered in each plot, and their average percentage cover may be observed in Table 3. Regarding the species origin (indigenous or exotic), the vascular flora present in these plots fall into five categories: Azores endemic (END), Macaronesia endemic (MAC), native non-endemic (N) (which, together, are the indigenous species), naturalized species (NATU) (including exotic and invasive species) and some species with doubtful colonization status (D) (Silva et al. 2010).

In order to characterize the vegetation communities a brief description of the 12 sampled sites is presented in the following section; the plots are illustrated on Figure 2 (sites 1–6) and Figure 3 (sites 7–12).

Site 1 (~10 m) — “Ponta da Ilha, Manhonha” — Coastal scrubland around the lighthouse of “Manhonha”, near “Piedade” village. A lava field divided by some stone walls, probably, the area was earlier used for orchards, which were abandoned from the middle of the 20th century. Presently, the plots are dominated by *Morella faya* which reached a cover of 75% (Table 3) and, at the herbaceous level, by *Pteridium aquilinum* (average about 60%). Among others, the endemic species *Erica azorica*, *Picconia azorica*, *Juniperus brevifolia*, *Euphorbia azorica* were also registered in the area. *M. faya* was the tallest tree registered on the plots, reached almost 6 m (height=580 cm; dbh=22 cm). The naturalized *Salpichroa organifolia* (Lam.) Baill. (Silva et al. 2008d), was registered for Plot 2, and is now documented in the entire archipelago except in Corvo Island (Silva et al. 2010).

Site 2 (~200 m) — “Cabeço da Hera” — lowland forest located near “Piedade” village. The surrounding area of “Cabeço da Hera” is occupied by a quarry, pastures for cattle, and farmland with some maize cultivation. Both plots

had a very steep slope, of about 50° (Table 2). More than a third of all species (N=9; 39%) were exotic, and *Pittosporum undulatum* exhibited the highest cover (Table 3); this is in line with the Disturbance Index, which presents the second highest value observed in the transect (D=47; Table 2). Nevertheless, the tallest trees recorded on site were the endemics *Picconia azorica* (height=600 cm; dbh=11 cm) and *Morella faya* (height=590 cm; dbh=20 cm).

Site 3 (~400 m) — “Fetais, Piedade” — forest. The plots are close to a secondary road, and grasslands, interspersed with forested hills covered with *Morella faya*, *Laurus azorica* and *Pittosporum undulatum* dominate the surrounding area. These plots proved to be the most affected by anthropogenic influence, presenting also the highest Disturbance Index of the transect (D=57; Table 2). In fact this site is dominated by the presence of *Pittosporum undulatum* (c. 80%) (Table 3) reaching more than 8 m height (dbh=18 cm), but endemic species such as *Picconia azorica*, *Laurus azorica*, *Erica azorica*, *Ilex perado* subsp. *azorica* and the orchid *Platanthera micrantha* (Hochst. ex Seub.) Schlecht. were also present. And for other plant groups, the very rare endemic moss *Echinodium renauldii* (Cardot) Broth. was also found in this site (Coelho et al. 2013a).

Site 4 (~600 m) — “Caminho dos Burros, Chão Verde” — *Laurus azorica* and *Ilex perado* subsp. *azorica* dominate the vegetation (c. 50%). A specimen of *Erica azorica* was the tallest tree in the plots (height=640 cm; dbh=12 cm) and a specimen of *Laurus azorica* presented the largest diameter (dbh=18 cm). These plots showed the lowest Disturbance Index (D=15) of the entire gradient (Table 2). With 42 plant species (four naturalized), this is the richest site of the elevational gradient. It is possible to observe 23 out of the 25 differential species of the alliance *Juniperion brevifolii* Sjögren (1973), including: the ferns *Culcita macrocarpa* C. Presl, *Dryopteris aemula* (Aiton) O. Kuntze, *Hymenophyllum tunbrigense* (L.) Sm; the shrubs *Vaccinium cylindraceum* Sm., *Myrsine africana*; and the plant rarities *Platanthera micrantha* and *Sanicula azorica* Guthn. ex Seub.

Vascular plants on Pico Island transect



Fig 2. Illustration of plots at different standard elevations in Pico Island (a) 10 m [P2]; b) 200 m [P1]; c) 400 m [entrance]; d) 600 m [P1]; e) 800 m [entrance]; f) 1000 m [P1]).




Fig. 3. Illustration of plots at different standard elevations in Pico Island. Pico Mountain: (a) 1200 m [P1]; b) 1400 m [P2]; c) 1600 m [P1]; d) 1800 m [P2]; e) 2000 m [P1]; f) 2200 m [P2]).

Division	Class	Order	Family	Species	Col.	0010P1	0010P2	0200P1	0200P2	0400P1	0400P2	0600P1	0600P2	0800P1	0800P2	1000P1	1000P2	1200P1	1200P2	1400P1	1400P2	1600P1	1600P2	1800P1	1800P2	2000P1	2000P2	2200P1	2200P2		
			Polypodiaceae	<i>Polypodium azoricum</i> (Vasc) R. Fern.	END	<1			<1	<1	<1	<1	1																		
			Thelypteridaceae	* <i>Christella dentata</i> (Forssk.) Brownsey & Jermy	NATU			1				4	9		<1	<1		<1						<1							
				<i>Oreopteris limbosperma</i> (All.) Holub	N					<1		<1	<1	<1																	
				<i>Stegnogramma pozoi</i> (Lag.) K. Iwats.	N			9																							
			Woodsiaceae	<i>Deparia petersenii</i> (Kunze) M. Kato	NATU							<1																			
				<i>Diplazium caudatum</i> (Cav.) Jermy	N				<1	19	<1																				
Pinophyta																															
Pinopsida																															
Pinales																															
			Cupressaceae	* <i>Juniperus brevifolia</i> (Seub.) Antoine	END	17	1					3	14	13	18	35	41	<1	11												
Magnoliophyta																															
Magnoliopsida																															
Laurales																															
			Lauraceae	<i>Laurus azorica</i> (Seub.) Franco	END					18	15	60	36	7	4	6	7														
Ranunculales																															
			Papaveraceae	<i>Fumaria muralis</i> Sonder ex Koch subsp. <i>muralis</i>	NATU		<1																								
Fagales																															
			Myricaceae	<i>Morella faya</i> (Aiton) Wilbur	N	75	76	41	37			<1	4																		
Rosales																															
			Rhamnaceae	<i>Frangula azorica</i> V. Grubov	END							9	2		1																
			Rosaceae	<i>Duchesnea indica</i> (Andr.) Focke	NATU													2						2	4						
				<i>Fragaria vesca</i> L.	D													1													
				<i>Potentilla anglica</i> Laich.	N													3	<1	10	20	<1	4								
				<i>Rubus hochstetterorum</i> Seub.	END							<1	3																		
				<i>Rubus ulmifolius</i> Schott	NATU				2	<1	2	1	4																		
Malpighiales																															
			Euphorbiaceae	<i>Euphorbia azorica</i> Seub.	END	2																									
				<i>Euphorbia stygiana</i> H. C. Watson subsp. <i>stygiana</i>	END									37	9																
			Hypericaceae	<i>Hypericum foliosum</i> Aiton	END				1			<1					1														
			Violaceae	<i>Viola odorata</i> L.	NATU														<1												
				<i>Viola palustris</i> L. subsp. <i>juresii</i> (Link ex K. Wein) Cout.	N														<1												
Malvales																															

Site 5 (~800 m) — “Caiado” — Forest dominated by *Ilex perado* subsp. *azorica* (Table 3) where it reaches the highest height and dbh values (height=480 cm; dbh=16 cm). Among the tree species there are also *Juniperus brevifolia* and *Laurus azorica* while *Euphorbia stygiana* is distinguished by its large size, reaching heights of 270 cm. The ferns *Dryopteris crispifolia* Rasbach, Reichstein & Vida, *D. azorica* (Christ) Alston (c. 40% cover) and *Culcita macrocarpa* are also quite frequent. Surrounding this area, there is a lagoon, “Lagoa do Caiado”, forest areas (production, protection, recreation) and low intensity pastures (DROTRH 2008).

Site 6 (~1000 m) — “Caveiro” — Mountain cloud forest; system dominated by *Ilex perado* subsp. *azorica* and *Juniperus brevifolia* (Table 3). *Ilex perado* subsp. *azorica* was the tallest tree (height=370 cm) and *Juniperus brevifolia* the broadest one (dbh=17 cm); both species had high percentage covers (average 50% and 38%, respectively). Many other plants are considered differential species in the alliance *Juniperion brevifolii* (Sjögren 1973). The fern *Culcita macrocarpa* reached the highest abundance (60% cover). This site is part of the Natural Reserve of “Caveiro”, which includes both forests and bogs, but the presence of cattle is notorious in the semi-natural pastures surrounding the plots.

Sites 7 to 9 (~1200 to ~1600 m) — Pico Mountain — from this point on, the transect follows the mountain trail, which starts around 1200 m and extends up to the crater of Pico Mountain. Plants reaching more than 2 m were not present in the trail. The first site (site 7) is covered by an *Erica* scrub (c. 50% cover) reaching up to 200 cm. The fern *Blechnum spicant* (L.) Sm. occupied about a third of the area of the plot but *Calluna vulgaris*, *Ilex perado* subsp. *azorica* and *Vaccinium cylindraceum* did not reach 15% of cover. At 1400 m (site 8), vegetation stature is lower, with the tallest species reaching up to 75 cm high; *Calluna vulgaris* dominates the vegetation (> 50% cover) while short shrubs of *Erica azorica* cover about a quarter of the area and the endemic *Holcus rigidus* about a tenth (Table 3). At 1600 m (site 9) *Calluna vulgaris* (c. 60% cover) and *Erica azorica* (c. 50% cover) predominated, the latter reaching a canopy’s maximum height of 120 cm.

Site 10 to 12 (~1800 to ~2200 m) — Pico Mountain — these last three sites are dominated mainly by *C. vulgaris*, but *Daboecia azorica* and *Thymus caespititius* (Table 3) are also present. These plots present a low number of species, and no naturalized plants were recorded. At 1800 m (site 10), the canopy’s maximum height was 29 cm, in the remaining elevations, the species were even shorter, reaching up to 12–22 cm height. The endemic species, *Silene uniflora* Roth subsp. *cratericola* (Franco) Franco described from the crater of the mountain was recorded at 2200 m (site 12).

PATTERNS OF RICHNESS AND DISTRIBUTION OF VASCULAR PLANTS

The twelve elevational sites surveyed resulted in a list of 88 taxa — one species new to the flora of the island and five identified only to the genus level — which represent circa 14% of all vascular taxa present in this island. A total of 60 indigenous species (32 END, 2 MAC and 26 N), 21 exotic species, two species with doubtful colonization status and five species identified to genera were found in the surveyed sites.

About a fifth of the species (N=20; 23%) presented a broad distribution range and were present in five or more elevations (Table 3, species marked with an asterisk): six ferns, the Azorean cedar, 10 Magnoliopsida, two Liliopsida and one Selaginellopsida. Except for the fern *Christella dentata*, all species were indigenous. Most of the species (N=51; 58%) occur up to 1000 m, while about a fifth (N=18; 21%) were only recorded above that altitude, including *Calluna vulgaris*, *Daboecia azorica*, *Thymus caespititius*, *Festuca francoi* Fern. Prieto, C. Aguiar, E. Dias & M. I. Gut and *Silene uniflora* subsp. *cratericola* (Table 3).

In the transect, eudicotyledons’ richness (Class Magnoliopsida) is irregularly distributed among the different elevations surveyed, ranging from three to 20 species, with the highest sites (1800 to 2200 m) showing fewer species (Table 3; Fig. 4). Maximum diversity is achieved by Magnoliopsida at 600 and 1200 m, with the presence of 20 and 18 species, respectively. Monocotyledons’ richness (Class Liliopsida) ranges between one to six species. Liliopsidas’ highest diversity is reached at 600 and 1000 m

Vascular plants on Pico Island transect

(Fig. 4). The only gymnosperm found in the transect, the endemic species *Juniperus brevifolia* (Division Pinophyta, Class Pinopsida), occurs in both plots of five sites (10, 600, 800, 1000 and 1200 m) (Table 3). Ferns' richness (Class Polypodiopsida) reaches 14 species at 600 m and 10 species at 200 and 800 m (Fig. 4). *Huperzia suberecta* and *Selaginella kraussiana* are the two representatives of the Classes Lycopodiopsida and Selaginellopsida, respectively; the first is present only at 1000 m while the second is common between 400 and 1200 m (Table 3; Fig. 4).

The number of indigenous species (END, MAC and N) was always higher than the number of exotic species (NATU). The maximum relative richness of indigenous species was recorded around the middle of the elevational gradient at 600, 800 and 1000 m (Fig. 5). These altitudes correspond to the most complex forests present in

the transect, including the trees *Laurus azorica*, *Ilex perado* subsp. *azorica* and *Juniperus brevifolia* and many other woody species. Exotic species richness reaches its maximum value at 10 and 200 m. At the remaining elevation levels the relative richness of exotic species is minimal, less than 2%, and they are completely absent in the highest elevations, between 1800 and 2200 m (Table 2; Fig. 5).

The distribution of species richness seems to follow a parabolic pattern (Fig. 4 and 5). Nevertheless, the indigenous vascular plant species distribution does not fit the Mid Domain Effect (MDE) (Fig. 6), since both the peak of richness is reached below the middle of the gradient and the number of species in the highest elevations is lower than expected. Similar results were obtained for all the species including exotics (not shown).




Fig. 4. Number of vascular plants per class at each studied standard elevation.




Fig. 5. Proportion of species: indigenous (END, endemic species to Azores; MAC, endemic species to Macaronesia; N, native species); exotic (NATU, naturalized species); and with doubtful colonization status (D) along the Pico transect.




Fig. 6. Mid Domain Effect (MDE) test for indigenous species distribution on the Pico Island transect.

DISCUSSION

Among the classification systems describing the Azorean vegetation, many used elevation to explain biodiversity patterns (see Table 1 and references therein). This is understandable since not only does biodiversity respond to elevation in a number of ways (Rahbek 1995) but, in general, it is also considered a good surrogate of habitat diversity and climatic variables, i.e. “environmental heterogeneity” (e.g. Hortal et al. 2009; Kessler et al. 2011; Ah-Peng et al. 2012; Gabriel et al. 2014). In the Azores, some authors have proposed exposure to winds and soil’s humidity as the main factors explaining the distribution of plant communities (e.g. Dias 1996; Dias et al. 2005), nevertheless, the elevational approach used in this study, analysing only a minimum fraction of the available area of the island (24 plots of 100 m²), resulted in a list of more than a third of the indigenous species (N=60; 35%), although intermixed with some of the exotic species (N=21; 5%) present in Pico Island. This transect and the permanent marked plots, thus represents a good start for future monitoring studies.

The composition found in the surveyed plots roughly agrees with the studies of Dansereau (1966), since this author places a prominent value on species such as *Morella faya* (0–300 m), *Laurus azorica* (300–650 m) and *Juniperus brevifolia* (450–1100 m). There are also striking resemblances with the Alliance *Juniperion brevifolii* described by Sjögren (1973), since a large number of differential species was also retrieved within the plots, especially above 400 m, where the proportion of species belonging to *J. brevifolii* and their average cover is always higher than the species with no differential value (data not shown).

The richness of species steadily rises from the coast (N=18: 11 indigenous and 7 exotic) until the 600 m (N=42: 38 indigenous and 4 exotic) where it peaks, declining afterwards to a very low number of species inside the caldera of Pico Mountain (Table 2; Fig. 5). Although the species distribution resembles a normal curve, it does not support the Mid Domain Effect hypothesis (Fig. 6), since the peak of richness appears much lower than the

middle of the domain (1200 m). A similar deviation from the predictions was also found with the epiphytic bryophytes of Réunion Island–Mascarene archipelago (Ah–Peng et al. 2012).

Many Azorean endemic plants are characterised by a wide ecological and elevational range (e.g. Sjögren 1973; Schäfer 2005; Table 1) and these modern times’ observations are also corroborated by palaeoclimatic studies (Connor et al. 2012, 2013). In fact, the Pico stratified sampling showed 20 species with ranges equal or higher than 1000 m (Table 3); the widest range belongs to *Erica azorica* (c. 1600 m range), a species considered as a cornerstone of the Azorean native vegetation (e.g. Ribeiro & Borges 2010). Other endemics species, such as *Juniperus brevifolia*, still present a wide range (c. 1200 m), but was missing from the sites of 200–400 m. This means that some species are becoming rarer, and that the indigenous communities are fragmented which agrees with the idea put forward by many authors (Table 1), for instance Purvis & James (1993: 2) who stated that in the Azores “virtually no natural forests with native trees exist below 500 m, having been replaced almost completely by the exotic *Pittosporum undulatum* (...)”. Indeed the sites surveyed at lower elevations (200 and 400 m), were impoverished and fragmented, dominated by *P. undulatum* (Table 3).

In fact, invasion pressure continues in the Azores, as is exemplified by the new record for Pico Island of the naturalized *Salpichroa origanifolia* (10 m site), a species considered among the Top 100 invasive species of Macaronesia (Silva et al. 2008d).

As expected (e.g. Silva et al. 2005) exotic species richness was higher in the sites where human action is dominant. Nevertheless, it was possible to find eight exotic plant species along the mountain trail (1200–1600 m), all presenting low covers (below 5% except *A. Odoratum* L., with 7%) (Table 3) and only one (*D. Indica* (Andr.) Focke) currently considered invasive (Fernandes et al. 2008). Although these species were introduced a long time ago in the Azores (cf. Schäfer 2003), and the nature of invasive species promote their installation, its presence above 1000 m could be exacerbated by the intensity of visitors arriving in

the lowland, ready to hike to the top of the mountain, stepping into the natural vegetation and accidentally transporting propagules of exotic species (it is possible to find seedlings or young trees of apple, peach and apricot along the trail up to the Pico summit, clearly indicating that visitors do have an effect in vegetation). Similarly to other touristic areas (e.g. Huiskes et al. 2014), this ecotourism pressure may promote the spread of additional invasive species to higher altitudes in Pico, calling to urgent measures of tourists self-regulation or special measures of education by the park rangers in order to reduce propagule transfer of plants.

Invasions and habitat loss (Martín et al. 2008) are serious threats to the indigenous flora (e.g. Schäfer 2005; Heleno 2008; Silva et al. 2008a, 2009), hindering both natural evolutionary processes and ecosystem services. Besides, changes in the dynamics of the vegetation from the Azores may be reinforced by the foreseen climatic changes (Harter et al. 2015).

Notwithstanding it is worth reinforcing that forests at medium–higher elevations (600–1000 m) harbour an impressive number of endemic vascular plant species (Table 2, Fig. 5) establishing native communities similar to the ones described in the past (e.g. Seubert & Hochstetter 1843; Morelet 1860; Tutin 1953; Dansereau 1966; Sjögren 1973; Dias 1996). This richness was also confirmed for other plant groups, such as bryophytes (Coelho et al. 2013b). At the lower altitude sites, and in spite of their obvious alterations, it was possible to list a relatively high number of woody endemic species along with *Platanthera micrantha* and endemic ferns, while the disturbance index of subalpine and alpine habitats remains quite low.

Currently, the native vegetation of Pico Island remains very rich and important for conservation and, although all sites but one (400 m) are well within the boundaries of the Pico Natural Park, it is necessary to monitor them to validate and/or adjust eventual conservation measures, such as the control of invasive species — PRECEFIAS Program for Control or Eradication of Invasive Plants in Sensitive Areas (Silva et al. 2008b) — or the restauration of habitats using local species to better guarantee the survival of populations and

preserve their evolutionary potential (e.g. Rumeu et al. 2011). Monitoring and management of these sites will be very important to prevent future threats to native areas. Furthermore the ongoing fragmentation (REAA 2005) and degradation of habitats due to trampling (e.g. Silva et al. 2005), expansion of exotic flora (e.g. Silva et al. 2008b) and replacement of indigenous vegetation by grasslands or plantation of non–endemic trees (e.g. Triantis et al. 2010) are threats to the Azorean biodiversity (Borges et al. 2009) that may change the quality of the studied ecosystems in the near future.

ACKNOWLEDGMENTS

The authors would like to acknowledge the financial support granted by the research project MOVECLIM (M2.1.2/F/04/2011/NET) and also by projects ISLANDBIODIV (NETBIOME/0003/2011), IMPACTBIO (DRCT–M2.1.2/I/005/2011) and ATLANTISMAR (DRCT–M2.1.2/I/027/2011). Besides, MCMC and DSGH were funded by Ph.D. grants M3.1.2/F/051/2011 and M3.1.2/F/051/2011, from the “Fundo Regional Ciência e Tecnologia”, Regional Government of the Azores and PROEMPREGO. The team are grateful to the Director of the Natural Park of Pico Island, Dr. Paulino Costa, and to the Director of the research group CITA–A — “Centro de Investigação e Tecnologias Agrárias–Açores”, Doutor João Madruga, for logistic assistance. The authors wish to thank Pedro Cardoso for estimating disturbance data for each site and the referees, Hanno Schäfer and Mónica Moura, for their valuable comments on an earlier version of the manuscript.

REFERENCES

- Ah–Peng, C., N. Wilding, J. Kluge, B. Descamps–Julien, J. Bardat, M. Chuah–Petiot, D. Strasberg et al. 2012. Bryophyte diversity and range size distribution along two altitudinal gradients: Continent vs. Island. *Acta Oecologica* 42: 58–65.
- Allorge, P. & V. Allorge 1946. Les étages de la végétation muscinale aux îles Açores et leurs éléments. *Mémoires de la Société de*

- Biogéographie* 8: 369–386. [The stages of the bryophytic vegetation in the Azores islands and their elements; in French]
- Archavaleta, M., S. Rodríguez, N. Zurita & A. García 2010. *Lista de Especies Silvestres de Canarias. Hongos, Plantas y Animales Terrestres (1a edición)*. Gobierno de Canarias. 579 pp. [List of Wild Species of Canarias. Fungi, Plants and Terrestrial Animals; in Spanish]
- Azevedo, E.B. 1996. Modelação do clima insular à escala local. Modelo CIELO aplicado à ilha Terceira. PhD Thesis, Universidade dos Açores. 247 pp. [Insular climate modeling at the local scale. CIELO model applied to Terceira Island; in Portuguese]
- Azevedo, E.B., L.S. Pereira & B. Itier 1999a. Simulation of local climate in islands environments using a GIS interated model. In: Musy A., L.S. Pereira & M. Fritsch (Eds.) *Emerging Technologies for Sustainable land Use and Water Management 2nd Inter-Regional Conference Environment-Water*, Lausanne Switzerland. CD-ROM Paper 7.4.
- Azevedo, E.B., L.S. Pereira & B. Itier 1999b. Modelling the local climate in island environments: water balance applications. *Agricultural Water Management* 40: 393–403.
- Azevedo, E.B., M.C. Rodrigues & J.F. Fernandes 2004. *Climatologia*. Pp. 25–48 in: Forjaz, V. H. (Eds). *Atlas Básico dos Açores*. Observatório Vulcanológico e Geotérmico dos Açores, Ponta Delgada. 112 pp. [Climatology; in Portuguese]
- Barceló, A.M. & L.F. Nunes 2011. *Atlas Climático dos Arquipélagos das Canárias, da Madeira e dos Açores*. Departamento de Produção da Agência Estatal de Meteorologia de Espanha e Departamento de Meteorologia e Clima do Instituto de Meteorologia de Portugal. 78 pp. [Climatic Atlas of the Canaries, Madeira and the Azores Archipelagos; in Portuguese]
- Borges, P.A.V., C. Abreu, A.M.F. Aguiar, P. Carvalho, S. Fontinha, R. Jardim, I. Melo, et al. 2008. Terrestrial and freshwater biodiversity of the Madeira and Selvagens archipelagos. Pp. 13–25 in: Borges, P.A.V., C. Abreu, A.M.F. Aguiar, P. Carvalho, R. Jardim, I. Melo, P. Oliveira, et al. (Eds). *A List of the Terrestrial Fungi, Flora and Fauna of Madeira and Selvagens Archipelagos*. Direcção Regional do Ambiente da Madeira e Universidade dos Açores, Funchal e Angra do Heroísmo. 440 pp.
- Borges, P.A.V., I.R. Amorim, R. Cunha, R. Gabriel, A.F. Martins, L. Silva, A. Costa & V. Vieira 2009. Azores. Pp 70–75 in: Gillespie, R.G. & D. Clagu (Eds). *Encyclopedia of Islands*. University of California Press, California. 1074 pp.
- Borges, P.A.V., J. Bried, A. Costa, R. Cunha, R. Gabriel, V. Gonçalves, A.F. Martins et al. 2010. Description of the terrestrial and marine Azorean biodiversity. Pp. 9–33 in: Borges, P.A.V., A. Costa, R. Cunha, R. Gabriel, V. Gonçalves, A.F. Martins, I. Melo et al. (Eds). *A List of the Terrestrial and Marine Biota from the Azores*. Princípiã, Cascais. 432 pp.
- Cardoso, P., F. Rigal, S. Fattorini, S. Terzopoulou & P.A.V. Borges 2013. Integrating landscape disturbance and indicator species in conservation studies. *PLoS ONE* 8: e63294.
- Coelho, M., R. Gabriel, D. Henriques & C. Ah-Peng 2013a. Espécies raras de briófitos ao longo do gradiente altitudinal de floresta nativa na ilha do Pico (Açores): o caso de *Echinodium renauldii* (Cardot) Broth.. Pp. 129–134 in: *Livro de Atas — Jornadas Ciência nos Açores. Que Futuro?*, São Miguel-Açores, Junho 2013. 157 pp. [Rare bryophyte species along an altitudinal gradient of native forest in Pico Island (Azores): the species *Echinodium renauldii*; in Portuguese]
- Coelho, M., R. Gabriel, D. Henriques & P.A.V. Borges 2013b. Distribuição de briófitos epífitos raros na floresta nativa da ilha do Pico: Resultados preliminares. Pp. 68–73 in: *Livro de Actas do I Congresso de Ciência e Desenvolvimento dos Açores: Crise, Território e Paisagem*. ACDA—Associação para a Ciência e Desenvolvimento dos Açores, Terceira-Açores, Julho 2013. 291pp. [Distribution of rare epiphytic bryophytes in native forest of Pico Island: preliminary results; in Portuguese]
- Colwell, R.K. & G.C. Hurtt 1994. Nonbiological Gradients in Species Richness and a Spurious Rapoport Effect. *The American Naturalist* 144: 570–595.
- Colwell, R.K. & D.C. Lees 2000. The mid-domain effect: geometric constraints on the geography of species richness. *Trends in Ecology & Evolution* 15: 70–76.
- Connor, S.E., J.F.N. van Leeuwen, T.M. Rittenour, W.O. van der Knaap, B. Ammann & S. Björck 2012. The ecological impact of oceanic island colonization — a palaeoecological perspective from the Azores. *Journal of Biogeography* 39: 1007–1023.
- Connor, S.E., W.O. van der Knaap, J.F.N. van Leeuwen & P. Kuneš 2013. Holocene palaeoclimate and palaeovegetation on the islands of Flores and Pico. Pp: 149–162 in: Fernández-Palacios, J.M., L. Nascimento, J.C. Hernández, S. Clemente, A. González & J.P. Díaz-González (Eds). *Climate change perspectives from the*

- Atlantic: Past, present and future*. Servicio de Publicaciones, Universidad de La Laguna. Tenerife, Canárias. 727 Pp.
- Cruz, J.V., R. Pereira, A. Moreira 2007. *Carta de Ocupação do Solo da Região Autónoma dos Açores*. Ed. Secretaria Regional do Ambiente e do Mar, Direcção Regional do Ordenamento do Território e dos Recursos Hídricos. 56pp. [Azorean Soil Occupation Chart; in Portuguese]
- Dansereau, P. 1966. Études Macaronésiennes III. La zonation altitudinale. *Naturaliste Canadien* 93: 779–795 [Macaronesian Studies III. The altitudinal zonation; in French].
- Dias, E. 1996. Vegetação natural dos Açores. Ecologia e sintaxonomia das florestas naturais. PhD Thesis, University of Azores. 302 pp. [Natural vegetation of the Azores; in Portuguese]
- Dias, E., R.B. Elias & V. Nunes 2004. Vegetation mapping and nature conservation: a case study in Terceira Island (Azores). *Biodiversity and Conservation* 13: 1519–1539.
- Dias E., C. Mendes, C. Melo, D. Pereira & R.B. Elias 2005. Azores central Islands vegetation and flora field guide. *Quercetea* 7: 123–173.
- DLR, Decreto Legislativo Regional n.º 20/2008/A de 9 de Julho. Diário da República, 1.ª série–N.º 131–9 de Julho de 2008.
- DROTRH 2008. Direcção Regional do Ordenamento do Território e Recursos Hídricos. *Carta de ocupação do solo da Região Autónoma dos Açores*. Secretaria Regional do Ambiente. Direcção Regional do Ordenamento do Território e dos Recursos Hídricos. Ponta Delgada. [Azorean Soil Occupation Chart; in Portuguese]
- Elias, R.B. 2001. *Sucessão primária em domas traquíticos*. MSc Thesis, University of Azores. 167 pp. [Primary succession on lava domes; in Portuguese]
- Elias, R.B. 2007. Ecologia das florestas de *Juniperus* dos Açores. PhD Thesis, University of Azores. 179 pp. [Ecology of Azorean *Juniperus* forests; in Portuguese]
- Elias, R.B., E. Dias & F. Pereira 2011. Disturbance, regeneration and the spatial pattern of tree species in Azorean mountain forests. *Community Ecology* 12: 23–30.
- Fernandes, F.M., E. Ojeda & L. Silva 2008. *Duchesnea indica* (Andr.) Focke. Pp. 490–492 in: Silva, L., E.L. Ojeda, & J.L. Rodriguez–Luengo (Eds). *Invasive Terrestrial Flora & Fauna of Macaronesia. TOP 100 in Azores, Madeira and Canaries*. ARENA, Ponta Delgada. 546 pp.
- Forjaz, V.H. 2004. *Atlas Básico dos Açores*. Observatório Vulcanológico e Geotérmico dos Açores. Ponta Delgada. 112 pp. [Basic Atlas of the Azores; in Portuguese]
- França, Z., J.V. Cruz, J.C. Nunes & V.H. Forjaz 2003. Geologia dos Açores: uma perspectiva actual. *Açoreana* 10: 11–140. [Azores Geology: an actual perspective; in Portuguese]
- França, Z., V.H. Forjaz, C.A. Ribeiro, A.M. Vaz, E. Ribeiro, E.B. Azevedo, J.M. Tavares & L.M. Almeida 2014. *Guia de História Natural da Ilha do Pico*. Observatório Vulcanológico e Geotérmico dos Açores. 400 pp. [Pico Island Natural History Handbook; in Portuguese]
- Frutuoso, G. 1998. *Saudades da Terra*: livro VI. Instituto Cultural de Ponta Delgada, Açores, Portugal. 165 pp. [Longing for the Land: book VI; Originally published in 1589; in Portuguese]
- Gabriel, R., M.M.C. Coelho, D.S.G. Henriques, P.A.V. Borges, R.B. Elias, J. Kluge & C. Ah–Peng 2014. Long–term monitoring across elevational gradients to assess ecological hypothesis: a description of standardized sampling methods in oceanic islands and first results. *Arquipelago*. Life and Marine Sciences 31: 45–67.
- Gaspar, C., K.J. Gaston, P.A.V. Borges & P. Cardoso 2011. Selection of priority areas for arthropod conservation in the Azores archipelago. *Journal of Insect Conservation* 15: 671–684.
- Guppy, H.B. 1917. *Plants, Seeds and Currents in the West Indies and the Azores*. Williams and Norgate. London. 531 pp.
- Harter D.E., S.D. Irl, B. Seo, M.J. Steinbauer, R.G. Gillespie, K.A. Triantis, J.M. Fernández–Palacios & C. Beierkuhnlein 2015. Impacts of global climate change on the floras of oceanic islands–projections, implications and current knowledge. *Perspectives in Plant Ecology, Evolution and Systematics* 17: 160–183.
- Heleno, R.H. 2008. The impact of alien plants on native biota in the Azores: a food web approach. PhD Thesis, Bristol: Faculty of Science, University of Bristol. 179 pp.
- Hortal J., K.A. Triantis, S. Meiri, E. Thébault & S. Sfenthourakis 2009. Island species richness increases with habitat diversity. *The American Naturalist* 174: E205–217.
- Hortal, J., P.A.V. Borges, A. Jiménez–Valverde, E.B. Azevedo & L. Silva 2010. Assessing the areas under risk of invasion within islands through potential distribution modelling: the case of *Pittosporum undulatum* in São Miguel, Azores. *Journal for Nature Conservation* 18: 247–257.
- Huiskes, A.H., N.J. Gremmen, D.M. Bergstrom, Y. Frenot, K.A. Hughes, S. Imura & S.L. Chown 2014. Aliens in Antarctica: Assessing transfer of plant propagules by human visitors to reduce invasion risk. *Biological Conservation* 171: 278–

- 284.
- INE (Internet). Instituto Nacional de Estatística. (cited March 2015). Available from: http://www.ine.pt/scripts/flex_definitivos/Main.html [Nacional Statistic Institute; in Portuguese]
- Kessler, M., J. Kluge, A. Hemp, & R. Ohlemüller 2011. A global comparative analysis of elevational species richness patterns of ferns. *Global Ecology and Biogeography* 20: 868–880.
- Leuschner, C. 1996. Timberline and alpine vegetation on the tropical and warm temperate oceanic islands of the world. *Vegetatio* 123: 193–206.
- Lima, E.A., J.C. Nunes, M.P. Costa & M. Machado 2014. Basis for the geological heritage management in the Azores Archipelago (Portugal). *Journal of Integrated Coastal Zone Management* 14: 301–319.
- Lüpnitz, D. 1975. Die vertikale Vegetationsgliederung auf der Insel Pico–Azoren. *Cuadernos de Botánica Canaria XXIII/XXIV*: 15–24. [The classification of the vertical vegetation of the Pico Island–Azores; in German]
- Madeira, M., J. Pinheiro, J. Madruga & F. Monteiro 2007. Soils of volcanic systems in Portugal. Pp. 69–81 in: Arnalds, O., F. Bartoli, P. Buurman, H. Oskarsson, G. Stoops & E. Garcia–Rodeja (Eds). *Soils of Volcanic Regions in Europe*. Springer–Verlag, Berlin. 645 pp.
- Marler, P. & D.J. Boatman 1952. An Analysis of the Vegetation of the Northern Slopes of Pico–The Azores. *Journal of Ecology* 40: 143–155.
- Martín, J.L., M. Arechavaleta, P.A.V. Borges & B. Faria (Eds). 2008. *TOP 100 As Cem Espécies Ameaçadas Prioritárias em Termos de Gestão na Região Europeia Biogeográfica da Macaronésia*. Consejería de Medio Ambiente y Ordenación Territorial, Gobierno de Canarias, 500 pp. [Top Hundred Priority Endangered Species in Managing Terms in the Biogeographical European Region of Macaronesia; in Portuguese]
- Menezes de Sequeira, M., D. Espírito–Santo, C. Aguiar, J. Capelo & J. Honrado 2012. *Checklist da Flora de Portugal (Continental, Açores e Madeira)*. Associação Lusitana de Fitossociologia. Lisboa, 74 pp. [Checklist of the Flora of Portugal (Mainland, Azores and Madeira); in Portuguese]
- Morelet, A. 1860. *Iles Açores*. Notice sur l'Histoire Naturelle des Açores Suivie d'une Description des Mollusques Terrestres de cet Archipel. J.–B. Bailliere et Fils, Paris, 216 pp. [Information on the Natural History of the Azores Followed by a Description of Terrestrial Molluscs of the Archipelago; in French]
- Nunes, J.C. 1999. A actividade vulcânica na ilha do Pico do Plistocénio Superior ao Holocénio: Mecanismo eruptivo e hazard vulcânico. PhD Thesis, University of Azores. 357 pp. [Volcanic activity in Pico Island from the Upper Pleistocene to the Holocene: eruptive mechanism and volcanic hazard; in Portuguese]
- PNA (internet). Parques Naturais dos Açores. (cited January 2016). Available from: <http://parquesnaturais.azores.gov.pt/pt/> [Azorean Natural Parks; in Portuguese]
- Purvis, O.W. & P.W. James 1993. Studies on the lichens of the Azores. Part 1 — Caldeira do Faial. *Arquipelago*. Life and Marine Sciences 11A: 1–15.
- Purvis, O.W., C.W. Smith & P.W. James 1994. Studies in the lichens of the Azores. Part 2—lichens of the upper slopes of Pico mountain. A comparison between the lichen floras of the Azores, Madeira and the Canary Islands at high altitudes. *Arquipelago*. Life and Marine Sciences 12A: 35–50.
- Purvis, O.W., James, P.W. & C.W. Smith 1995. Studies on the lichens of the Azores. Part 3 — Macrolichens of relict cloud forests. *Boletim do Museu Municipal do Funchal* 4: 599–619.
- Rahbek, C. 1995. The elevational gradient of species richness: a uniform pattern? *Ecography* 18: 200–205.
- REAA 2005. *Relatório do estado do ambiente dos Açores 2005*. Secretaria Regional do Ambiente e do Mar. [Status Report of the Azorean Environment; in Portuguese]
- REAA 2014. *Relatório do estado do ambiente dos Açores 2011–2013*. Secretaria Regional da Agricultura e Ambiente, Direcção Regional do Ambiente. [Status Report of the Azorean Environment; in Portuguese]
- Ribeiro, S.P. & P.A.V. Borges 2010. Canopy habitat area effect on the arthropod species densities in the Azores: pondering the contribution of tourist species and other life histories. Pp. 89–114 in: Serrano, A.R.M., P.A.V. Borges, M. Boieiro & P. Oromí (Eds.). *Terrestrial Arthropods of Macaronesia — Biodiversity, Ecology and Evolution*. Sociedade Portuguesa de Entomologia, Lisboa. 327 pp.
- Rijsdijk, K.F., T. Hengl, S. Norder, R. Otto, B.C. Emerson, S.P. Ávila, H. López, E. van Loon, E. Tjorve & J.M. Fernández–Palacios 2014. Quantifying surface area changes of volcanic islands driven by Pleistocene sea level cycles: biogeographic implications for Macaronesian archipelagos, Atlantic Ocean. *Journal of Biogeography* 41: 1227–1439.
- Rumeu, B., J. Caujapé–Castells, J.L. Blanco–Pastor, R. Jaén–Molina, M. Nogales, R.B. Elias & P. Vargas 2011. The colonization history of *Juniperus*

- brevifolia* (Cupressaceae) in the Azores Islands. *PLoS ONE* 6: e27697.
- Schäfer, H. 2003. Chorology and diversity of the Azorean flora. *Dissertationes Botanicae* 374: 1–130.
- Schäfer, H. 2005. Endemic vascular plants of the Azores: an updated list. *Hoppea–Denkschriften der Regensburgischen Botanischen Gesellschaft* 66: 275–283.
- Schäfer, H., M. Moura, M.G.B. Maciel, L. Silva, F. Rumsey & M.A. Carine 2011 The Linnean shortfall in oceanic island biogeography: a case study in the Azores. *Journal of Biogeography* 38: 1345–1355
- Seubert, M. & C. Hochstetter 1843. Übersicht der Flora der azorischen Inseln. *Archiv für Naturgeschichte* 9: 1–24. [Overview of the Flora of the Azorean Islands; in German]
- Silva, L., N. Cordeiro, X. Illas & A. Martinez 2005. Amostragem altitudinal da flora vascular e da vegetação na montanha do Pico. XII Expedição Científica do Departamento de Biologia — Universidade dos Açores. *Relatórios e Comunicações do Departamento de Biologia* 34: 212–218. [Altitudinal sampling of the vascular flora and vegetation of Pico Mountain; in Portuguese]
- Silva, L., E. Ojeda Land & J.L. Rodríguez–Luengo (Eds) 2008a. *Invasive Terrestrial Flora & Fauna of Macaronesia. TOP 100 in Azores, Madeira and Canaries*. ARENA, Ponta Delgada. 546 pp.
- Silva, L., E. Ojeda Land, J.L. Rodríguez–Luengo, P.A.V. Borges, P. Oliveira & R. Jardim 2008b. Invasive alien species in Macaronesia. Pp. 159–165 in: Silva, L., E. Ojeda Land, & J.L. Rodríguez–Luengo (Eds) *Invasive Terrestrial Flora & Fauna of Macaronesia. TOP 100 in Azores, Madeira and Canaries*. ARENA, Ponta Delgada. 546 pp.
- Silva, L., R. Corvelo, M. Moura, R. Mesa Coello & J.A. Carvalho 2008c. *Pittosporum undulatum* Vent. Pp. 225–228 in: Silva, L., E. Ojeda Land, & J.L. Rodríguez–Luengo (Eds.) *Invasive Terrestrial Flora & Fauna of Macaronesia. TOP 100 in Azores, Madeira and Canaries*. ARENA, Ponta Delgada. 546 pp.
- Silva, L., R. Corvelo, M. Moura, R. Mesa Coello & J.A. Carvalho 2008d. *Salpichroa origanifolia* (Lam.) Thell. Pp. 233–235 in: Silva, L., E.L. Ojeda, & J.L. Rodríguez–Luengo (Eds). *Invasive Terrestrial Flora & Fauna of Macaronesia. TOP 100 in Azores, Madeira and Canaries*. ARENA, Ponta Delgada. 546 pp.
- Silva, L., M. Martins, M. Moura & G.B. Maciel 2009. *Azorean Vascular Flora: Priorities in Conservation*. Amigos dos Açores & CCPA, Ponta Delgada. 116 pp.
- Silva, L., M. Moura, H. Schäfer, F. Rumsey, E.F. Dias 2010. List of Vascular Plants (Tracheobionta). Pp. 117–146 in: Borges, P.A.V., A. Costa, R. Cunha, R. Gabriel, V. Gonçalves, A. F. Martins, I. Melo, et al. *A List of the Terrestrial and Marine Biota from the Azores*. Princípiã, Cascais. 432 pp.
- Silva, L. 2015. How ecotourism works at the community–level: the case of whale–watching in the Azores. *Current Issues in Tourism* 18: 196–211.
- Silveira, L. 2013. *Aprender com a História — Modos de Interação com a Natureza na Ilha Terceira (Do Povoamento ao século XX)*. Princípiã. 357 pp. [Learning from History — Ways of interacting with Nature in Terceira Island (From settlement to the 20th century; in Portuguese)]
- Sjögren, E. 1973. Recent changes in the vascular flora and vegetation of the Azores Islands. *Memórias da Sociedade Broteriana* 22: 1–453.
- Sjögren, E. 1978. Bryophyte vegetation in the Azores islands. *Memórias da Sociedade Broteriana* 26: 1–273.
- Triantis, K.A., P.A.V. Borges, R.J. Ladle, J. Hortal, P. Cardoso, C. Gaspar, F. Dinis, et al. 2010. Extinction debt on oceanic islands. *Ecography* 33: 285–294.
- Tutin, T.G. 1953. The vegetation of the Azores. *Journal of Ecology* 41: 53–61.
- Zbyszewski, G., C.R. Ferreira & O.V. Ferreira 1963. *Carta Geológica de Portugal — Ilha do Pico (Açores), Notícia Explicativa*. Serviços Geológicos de Portugal. Lisboa. [Geological map of Portugal — Pico Island (Azores), explanatory notes; in Portuguese]

Received 13 Nov 2015, Accepted 19 Feb 2016,
Published online 28 July 2016